

---

# EFIMOV TRIMERS UNDER STRONG CONFINEMENT

Pietro Massignan  
(Institute of Photonic Sciences, Barcelona)

---



in collaboration with  
Jesper Levinsen (Aarhus Institute of Advanced Studies)  
Meera Parish (University College London)









# STRONG CONFINEMENT EFFECTS

The dimensionality of the embedding space profoundly affects the system properties.

Examples:

- Anderson localization
- condensation & superfluidity





---

# OUTLINE

---

- Three identical bosons: 3D vs. 2D
- What happens in between? (quasi-2D)
  - trimer spectra and aspect-ratios
  - hyper-spherical potentials and wave functions
- Experimental consequences




# 2&3 IDENTICAL BOSONONS IN 3D

One universal dimer:  $E_b = -\frac{\hbar^2}{ma^2} \quad (a > 0)$


For resonant interactions ( $1/a=0$ ), in principle  $\exists$  an infinite tower of Efimov trimers.

Trimers map onto each other via the scale transformations  $a \rightarrow \lambda_0^n a$  and  $E \rightarrow \lambda_0^{-2n} E$

$\lambda_0 = 22.7$



Braaten & Hammer, Phys. Rep. 2006



Largest set by the temperature, or the dimension of the container

Scaling symmetry: continuous (two-body) vs. discrete (three-body)




# 2&3 IDENTICAL BOSONS IN 2D

Apply harmonic confinement:  $V(z) = \frac{1}{2}m\omega_z^2 z^2$

- CoM decouples
- continuum is shifted
- additional length scale appears

$$l_z = \sqrt{\frac{\hbar}{m\omega_z}}$$



One universal dimer:  $E_b = \frac{\hbar\omega}{2} - \frac{\hbar^2}{ma_{2D}^2}$


Petrov & Shlyapnikov PRA 2001  
Bloch, Dalibard, Zwerger RMP 2008

Two universal trimers:

$$-1.27 \frac{\hbar^2}{ma_{2D}^2}$$

$$-16.5 \frac{\hbar^2}{ma_{2D}^2}$$

Bruch & Tjon, PRA 1979



Both two- and three-body problems display a continuous scaling symmetry




# THREE BOSONONS IN QUASI-2D

$$H = \sum_{\mathbf{k}, n} (\epsilon_{\mathbf{k}} + n\hbar\omega_z) a_{\mathbf{k}, n}^\dagger a_{\mathbf{k}, n} + \sum_{\substack{\mathbf{k}, \mathbf{k}', \mathbf{q} \\ n_1, n_2, n_3, n_4}} e^{-(\mathbf{k}^2 + \mathbf{k}'^2)/\Lambda^2} \langle n_1 n_2 | \hat{g} | n_3 n_4 \rangle a_{\mathbf{q}/2 + \mathbf{k}, n_1}^\dagger a_{\mathbf{q}/2 - \mathbf{k}, n_2}^\dagger a_{\mathbf{q}/2 - \mathbf{k}', n_3} a_{\mathbf{q}/2 + \mathbf{k}', n_4}$$

the UV cut-off  $\Lambda$  controls the three-body physics at short-distances, and fixes the crossing of the deepest Efimov trimer with the 3-atom continuum (a.)

Trimer wave function: 
$$\sum_{\substack{\mathbf{k}_1, \mathbf{k}_2 \\ n_1, n_2, n_3}} \psi_{\mathbf{k}_1, \mathbf{k}_2}^{n_1, n_2, n_3} a_{\mathbf{k}_1, n_1}^\dagger a_{\mathbf{k}_2, n_2}^\dagger a_{-\mathbf{k}_1 - \mathbf{k}_2, n_3}$$




J. Levinsen, P. Massignan, and M. Parish, arXiv:1402.1859



# SKORNIIAKOV—TER-MARTIROSIAN EQ.

atom-dimer vertex:



relative z-motion  
wave function at  $z_r=0$

Clebsch-Gordan coefficient

$$\mathcal{T}^{-1}(\mathbf{k}_1, E_3 - \epsilon_{\mathbf{k}_1} - N_1\omega_z) \chi_{\mathbf{k}_1}^{N_1} = 2 \sum_{\mathbf{k}_2, N_2, n_{23}, n_{31}} \frac{f_{n_{23}} f_{n_{31}} \langle N_1 n_{23} | N_2 n_{31} \rangle e^{-(k_1^2 + k_2^2)/\Lambda^2} \chi_{\mathbf{k}_2}^{N_2}}{E_3 - \epsilon_{\mathbf{k}_1} - \epsilon_{\mathbf{k}_2} - \epsilon_{\mathbf{k}_1 + \mathbf{k}_2} - (N_1 + n_{23})\omega_z}$$

(the CoM q.number  $N$  does not appear in the final formula!)

where:

$$\mathcal{T}(\mathbf{k}, E) = \frac{2\sqrt{2\pi}}{m} \left\{ \frac{l_z}{a} - \mathcal{F} \left( \frac{-E + k^2/4m}{\omega_\perp} \right) \right\}^{-1}$$

- $E_3$  is the energy measured from the 3-atom continuum
- $\mathbf{k}_i$  is the relative momentum of atom  $i$  w.r.t. the pair  $(j, k)$
- $n_{ij}$  and  $N_i$  are the h.o. quantum numbers for motion along  $z$  of a pair, and of an atom and a pair


Wave function for the  
atom-pair relative motion:  $\psi(\boldsymbol{\rho}, Z) = R^{3/2} \sum_{\mathbf{k}, N} e^{i\mathbf{k} \cdot \boldsymbol{\rho}} \phi_N(Z) \chi_{\mathbf{k}}^N$




# SPECTRUM

interaction strength:  $|a_-|/a$   
 confinement strength:  $\mathcal{C}_z \equiv |a_-|/l_z$

“weak” confinement



strong confinement



$^{133}\text{Cs}$ :  $\omega_z \approx 2\pi \times 5\text{kHz}$


$\omega_z \approx 2\pi \times 30\text{kHz}$

- deepest trimer closely resembles the 3D-one, even for strong confinement
- spectrum of trimers is strongly modified above the 3D continuum
- energy of trimer (measured from the q2D dimer) can be a significant fraction of  $\omega_z$  even when  $|a_-|/a < -1$ , so trimers can be quite *resistant to thermal dissociation* when  $T \ll \omega_z$




# SPECTRUM (2D STYLE)

$$C_z \equiv |a_-|/l_z$$



$^{133}\text{Cs}$ :  $\omega_z \approx 2\pi \times 5\text{kHz}$




$\omega_z \approx 2\pi \times 30\text{kHz}$

- the 2D limit is recovered for small and negative scattering lengths (“BCS side” of the resonance)
- the two deepest trimers are stabilized for every negative scattering length
- avoided crossings: superposition of trimers with Efimovian + 2D-like character



# SHAPE OF THE TRIMERS



2D limit

3D regime



---

# HYPERSPHERICAL POTENTIALS

---

$$R^2 = r_1^2 + r_2^2 + r_3^2$$

Hyper-spherical expansion:  $\Psi(R, \Omega) = \frac{1}{R^{5/2} \sin(2\alpha_k)} \sum_{n=0}^{\infty} f_n(R) \Phi_n(R, \Omega)$


Hyper-radial Schrödinger equation:

$$\left[ -\frac{1}{2m} \frac{\partial^2}{\partial R^2} + V(R) \right] f_0(R) = (E_3 + \omega_z) f_0(R)$$

$V(R)$  depends on  $l_z/a$ , but not on the 3-body parameter.



# HYPERSPHERICAL POTENTIALS




- $V(R)$  approaches the 3D potential for  $R \ll |a|$   
and the 2D potential for  $R \gg l_z$
- When  $l_z/a \lesssim -2.5$  the potential displays a repulsive barrier with height  $\sim 0.15/ma^2$
- Small weight of trimers in the short distance region enhances lifetime



# EXPERIMENTAL CONSEQUENCES

- As “2D” experiments are performed at confinements often weaker than 5kHz, we expect this crossover physics to impact three-body correlations in realistic 2D studies on the *attractive* side of the Feshbach resonance
- Confinement raises continuum by  $\hbar\omega_z$ , so trimer resonance and loss peak disappear for  $l_z/|a_-| \lesssim 2.5$ , i.e.,  $C_z \gtrsim 0.4$
- When aiming at observing the discrete scaling symmetry: the 2nd trimer signature disappears once  $C_z \gtrsim 0.4/22.7$  which for  $^{133}\text{Cs}$  corresponds to  $\omega_z \approx 2\pi \times 10\text{Hz}$
- Similar effects expected for 4-body states (as two tetramers exist in 2D), or in quasi-1D





---

# CONCLUSIONS

---

Efimov trimers under strong confinement


Discrete scaling survives only for  $|a_-| \ll |a| \ll l_z$

Deepest trimer remains 3D-like even under strong confinement

Mixing with 2D trimers stabilizes the two deepest trimers for all  $a < 0$

Small weight at short distance will enhance lifetime (long-lived Efimov trimers?)

Consequences for correlations, quest to observe discrete scaling symmetry







Thanks to:

Vudtiwat Ngampruetikorn (University of Cambridge)

Rudi Grimm, Francesca Ferlaino, Bo Huang & all the LevT team (Innsbruck)

J. Levinsen,  
P. Massignan and  
M. M. Parish,  
[arXiv:1402.1859](https://arxiv.org/abs/1402.1859)

And thank you all for the attention!